

Synthetic 3D inversion of tidally induced magnetic field

Libor Šachl, Jakub Velímský

6.1. 2021

Current section

- 1 Introduction
- 2 Forward and inverse modelling
- 3 Numerical setup
- 4 Results

Wind and buoyancy driven ocean circulation

- driven by wind stresses and pressure gradients
- **surface** and deep (= meridional overturning = thermohaline) circulation
- average horizontal velocities 0.01 - 0.1 m/s




Figure: Surface circulation (Garrison, 2012).

- driven by wind stresses and pressure gradients
- surface and **deep** (= meridional overturning = thermohaline) circulation
- average horizontal velocities 0.01 - 0.1 m/s




Figure: Deep circulation (Broecker et al., 1991; Schmitz Jr, 1996).

Previous seminars

- Baroclinic ocean model LSOMG (1.3. 2017)
- Elektromagnetická indukce v oceánech (25.4. 2017)
- Modelling of electromagnetic signatures of global ocean circulation: Physical approximations and numerical issues (27.2. 2019)
- EM induction in the Earth's Oceans II: Going big! (30. 10. 2019)

Today we focus on tides!

Tidal ocean circulation

- driven by gravitational forces of the Moon and the Sun




Figure: Tides caused by the Moon and the Sun
(<https://www.clipart.email>).

Tidal ocean circulation

- distinct tidal frequencies $\Rightarrow$ frequency-domain modelling


Name of tide	Symb	Period	Freq. (s^{-1})- ω	K (m)	Ampl.
Semidiurnal Species					
Principal Lunar	M_2	12.421h	$1.40519 \cdot 10^{-4}$	0.242334	
Principal Solar	S_2	12.000h	$1.45444 \cdot 10^{-4}$	0.112841	
Elliptical Lunar	N_2	12.658h	$1.37880 \cdot 10^{-4}$	0.046398	
Declination Solar	Luni- K_2	11.967h	$1.45842 \cdot 10^{-4}$	0.030704	
Diurnal Species					
Declination Solar	Luni- K_1	23.934h	$0.72921 \cdot 10^{-4}$	0.141565	
Principal lunar	O_1	25.819h	$0.67598 \cdot 10^{-4}$	0.100574	
Principal solar	P_1	24.066h	$0.72523 \cdot 10^{-4}$	0.046843	
Elliptical lunar	Q_1	26.868h	$0.64959 \cdot 10^{-4}$	0.019256	
Long-Period Species					
Fortnightly Lunar	Mf	13.661d	$0.053234 \cdot 10^{-4}$	0.041742	
Monthly Lunar	Mm	27.555d	$0.026392 \cdot 10^{-4}$	0.022026	
Semiannual Solar	Ssa	182.621d	$0.0038921 \cdot 10^{-4}$	0.019446	

Figure: Parameters of major tidal constituents (Kowalik et al., 2006)

Tidal ocean circulation

- gravitational force is almost constant within the water column
⇒ 2D flow

2020/03/04 00:12:00



Why tides?

- sensitive to upper mantle structure
 - suitable frequencies
 - quasi-global coverage
 - non-trivial structure
- CHAMP (10 years) + Swarm (5 years) $\Rightarrow$
 - M_2 ($j \leq 28$)
 - N_2 ($j \leq 12$)
 - O_1 ($j \leq 12$)

(Grayver and Olsen, 2019)
- 1 of 4 Swarm's key objectives: determine the 3D electrical conductivity of the mantle

Why tides?




Figure: Combined inversion from magnetospheric and M_2 data (Grayver et al., 2017)

Current section

1 Introduction

2 Forward and inverse modelling

3 Numerical setup

4 Results




Figure: Illustration of EM induction in the Earth's oceans and mantle.
Courtesy of Jan Dostál.

Governing equations

- quasi-static Maxwell equations supplemented by Ohm's law

$$\nabla \cdot \mathbf{B} = 0 \quad (1)$$

$$\nabla \times \mathbf{B} = \mu_0 (\mathbf{j} + \mathbf{j}^{\text{imp}}) \quad (2)$$

$$\nabla \times \mathbf{E} = -i\omega \mathbf{B} \quad (3)$$

$$\mathbf{j} = \sigma \mathbf{E} \quad (4)$$

- imposed electric field/currents

$$\mathbf{j}^{\text{imp}} = \sigma \mathbf{E}^{\text{imp}} = \sigma (\mathbf{u} \times \mathbf{B}_M)$$

- 3 assumptions:

$$\mathbf{B}_M = -\nabla V$$

$$\left| \frac{\partial \mathbf{B}_M}{\partial t} \right| \ll \left| \frac{\partial \mathbf{B}}{\partial t} \right|$$

$$|\mathbf{B}| \ll |\mathbf{B}_M|$$

ElmgFD

- solves the EM induction equation in the frequency domain

$$\nabla \times \left(\frac{1}{\sigma} \nabla \times \mathbf{B} \right) - i\omega\mu_0 \mathbf{B} = \mu_0 \nabla \times \mathbf{E}^{\text{imp}}$$

- complex vector spherical harmonic functions in angular coordinates
 - separation of poloidal and toroidal magnetic field modes
 - high contrast of electrical conductivity $\Rightarrow$ Gibbs phenomenon
 - difficult to locally refine the grid
- piecewise-linear 1-D finite elements in radial coordinate
- divergence-free constraint enforced by means of Lagrange multipliers
- solves a system of linear equations with block-tridiagonal matrix using iterative solver with a 1D preconditioner

Inversion

- model parameters m_i , $i \in \{1, \dots, M\}$

$$\log_{10} [\sigma(\mathbf{m}; \mathbf{r}) / \sigma_0] = \sum_{i=1}^M m_i \zeta_i(\mathbf{r}) + \sum_{i=M+1}^{M_0} m_i^0 \zeta_i(\mathbf{r})$$

- $\zeta_i(\mathbf{r})$ are piecewise-constant functions
- model parameters m_i^0 have fixed values (regions below continents)
- minimize penalty function (Velímský, 2013)

$$\tilde{\mathbf{m}}(\lambda) = \arg \min_{\mathbf{m} \in R^M} [\chi^2(\mathbf{m}) + \lambda R(\mathbf{m})]$$

$$\chi^2(\mathbf{m}) = \sum_l \sum_{jm} \frac{(2j+1)(j+1)}{8\pi} \left| \frac{G_{jm}^{(i)}(\mathbf{m}; \omega_l) - G_{jm}^{(i, \text{obs})}(\omega_l)}{\delta_{jm}^{(i)}(\omega_l)} \right|^2$$

Adjoint approach

We follow Maksimov and Velímský (2017).

Forward problem for ω_I :

$$\nabla \times [\rho(\mathbf{r}) \nabla \times \mathbf{B}(\mathbf{r}; \omega_I)] - i\omega_I \mu_0 \mathbf{B}(\mathbf{r}; \omega_I) = \mu_0 \nabla \times (\mathbf{v} \times \mathbf{B}_0)$$

Adjoint problem for ω_I :

$$\nabla \times [\rho(\mathbf{r}) \nabla \times \hat{\mathbf{B}}(\mathbf{r}; \omega_I)] + i\omega_I \mu_0 \hat{\mathbf{B}}(\mathbf{r}; \omega_I) = \mathbf{0}$$

or

$$\nabla \times [\rho(\mathbf{r}) \nabla \times \overline{\hat{\mathbf{B}}}(\mathbf{r}; \omega_I)] - i\omega_I \mu_0 \overline{\hat{\mathbf{B}}}(\mathbf{r}; \omega_I) = \mathbf{0}$$

Adjoint approach

Adjoint source for ω_l :

$$\hat{G}_{jm}^{(e,obs)}(\omega_l) = +i \frac{j+1}{2} \frac{G_{jm}^{(i)}(\mathbf{m}; \omega_l) - G_{jm}^{(i,obs)}(\omega_l)}{[\delta_{jm}^{(i)}(\omega_l)]^2}$$

or

$$\overline{\hat{G}_{jm}^{(e,obs)}}(\omega_l) = -i \frac{j+1}{2} \frac{\overline{G_{jm}^{(i)}(\mathbf{m}; \omega_l)} - \overline{G_{jm}^{(i,obs)}(\omega_l)}}{[\delta_{jm}^{(i)}(\omega_l)]^2}$$

Adjoint approach

Derivative of misfit with respect to m_α :

$$D_\alpha \chi^2(\mathbf{m}) = \frac{2}{4\pi a^3 \mu_0} \operatorname{Re} \left(\sum_l \frac{1}{\omega_l} \int_G l \, dV \right),$$

$$l = D_\alpha \rho(\mathbf{m}; \mathbf{r}) \nabla \times \mathbf{B}(\mathbf{m}; \mathbf{r}; \omega_l) \cdot \nabla \times \overline{\hat{\mathbf{B}}}(\mathbf{m}; \mathbf{r}; \omega_l),$$

where

$$D_\alpha \rho = -(\ln 10) \rho(\mathbf{m}; \mathbf{r}) \zeta_\alpha(\mathbf{r})$$

Current section

- 1 Introduction
- 2 Forward and inverse modelling
- 3 Numerical setup
- 4 Results

Forcing

- 2D TPX09 M_2 tidal velocities
 - $(1/30)^\circ \times (1/30)^\circ$
 - bilinearly interpolated to ElmgFD grid




Figure: Real (left) and imaginary (right) parts of zonal barotropic transports [m^2/s].

- 12th generation of the International Geomagnetic Reference Field (IGRF–12, Thébault et al., 2015) $\mathbf{B}_M$
 - constant with depth

Surface conductivity, ocean σ_{oc}

- mean σ_{oc} is 3.31 ± 0.23 S/m (3.2 S/m in our benchmarks)




Figure: Vertically averaged conductivity of the global ocean (Tyler et al., 2017).

Surface conductivity, ocean σ_{oc}

- we use σ_{oc} of Tyler et al. (2017)
 - annual means $\Rightarrow$ constant in time
 - depends on S , p and $T \Rightarrow$ spatially variable
 - 102 layers layers
 - from surface down to 5500 m




Figure: Surface conductivity (left) and globally averaged ocean conductivity profile (right) (Tyler et al., 2017).

Surface conductivity, sediments

Type	σ [S/m]
oceanic s. A	0.8
oceanic s. B	0.02
coastal s.	0.5
continental s.	0.03
igneous	0.001

Table: Conductivity of igneous rocks and sediments.




Table: Schematic layout of conductivity layers below continents, oceanic shelves and deep ocean.

3D conductivity of upper mantle, model WINTERC-e

Original model (Fullea et al., 2019):

- 0-402 km
- 201 layers, each 2 km thick
- irregular grid

We use averaged version on 90×180 Gauss-Legendre grid:

- vertical averaging: 10 layers
- horizontal averaging: 4-7 neighbouring data points




Figure: 1D profiles.

1D conductivity model of lower mantle and core




Figure: 1D conductivity models.

ElmgFD configuration

- spherical-harmonic truncation degree $j_{max} = 60$
- 90×180 ($2^\circ \times 2^\circ$) Gauss-Legendre grid
- total of 165 layers
 - 3D ocean layers: 102
 - 3D upper-mantle layers: 21
 - 1D core and lower-mantle layers: 43
- inverse problem
 - $6^\circ \times 6^\circ$ grid cells
 - 15 210 model parameters
 - 13 layers

Current section

- 1 Introduction
- 2 Forward and inverse modelling
- 3 Numerical setup
- 4 Results**




Figure: Target and inverted conductivity [S/m] in the depth of 153 km and 395 km. Internal coefficients on surface. No regularization.

OIMF in the outer space

$$\begin{aligned}\nabla \times \mathbf{B} &= 0 \quad \Rightarrow \quad \mathbf{B} = -\nabla U, \\ \nabla \cdot \mathbf{B} &= 0 \quad \Rightarrow \quad \Delta U = 0, \quad \lim_{r \rightarrow \infty} U = 0.\end{aligned}$$

$$U(\mathbf{r}, t) = a \sum_{j=1}^{\infty} \sum_{m=-j}^j G_{jm}^{(i)}(t) \left(\frac{a}{r}\right)^{j+1} Y_{jm}(\vartheta, \varphi)$$




Figure: Effect of height where internal coefficients are “measured” on the conductivity [S/m] determined by inversion. No regularization.




Figure: Effect of 1D/3D continental conductivity in the generation of synthetic internal coefficients on the conductivity [S/m] determined by inversion. No regularization.




Figure: Effect of maximum degree j_{maxD} of synthetic internal coefficients on the conductivity [S/m] in the depth of 153 km. No regularization.




Figure: Conductivity [S/m] in layer 56 in the presence of 5% noise for $j > 30$. Regularization: Horizontal derivative of logarithm of resistivity in all inspected layers.




Figure: Conductivity [S/m] in layer 56 in the presence of 5% noise for $j > 20$. Regularization: Horizontal derivative of logarithm of resistivity in all inspected layers.




Figure: Conductivity [S/m] in layer 56 in the presence of 5% noise for $j > 10$. Regularization: Horizontal derivative of logarithm of resistivity in all inspected layers.




Figure: Conductivity [S/m] in layer 56 in the presence of 1% noise for all j . Regularization: Horizontal derivative of logarithm of resistivity in all inspected layers.




Figure: Conductivity [S/m] in layer 56 in the presence of 3% noise for all j . Regularization: Horizontal derivative of logarithm of resistivity in all inspected layers.




Figure: Target, old and new inverted conductivity [S/m] in the depth of 153 km and 395 km. Internal coefficients at satellite height. No regularization.




Figure: Power spectra from data and our numerical simulations with 3D forcing.




Figure: Power spectra from data and our numerical simulations with 2D forcing.




Figure: Power spectra from data and our new numerical simulations with 3D forcing.

Summary

- we established a framework for the 3D inversion of tidally induced magnetic field
- synthetic tests are promising for the inversion of real data ...
- ... but the discrepancies between available data sets are discouraging

- Broecker, W.S., et al., 1991. The great ocean conveyor. *Oceanography* 4, 79–89.
- Fullea, J., Lebedev, S., Martinec, Z., Celli, N.L., 2019. WINTERC-grav: mapping the upper mantle thermochemical heterogeneity from coupled geophysical-petrological inversion of seismic waveforms. *Geophys. J. Int.*, under review.
- Garrison, T.S., 2012. *Essentials of oceanography*. Cengage Learning.
- Grayver, A., Munch, F., Kuvshinov, A., Khan, A., Sabaka, T., Tøffner-Clausen, L., 2017. Joint inversion of satellite-detected tidal and magnetospheric signals constrains electrical conductivity and water content of the upper mantle and transition zone. *Geophys. Res. Lett.* 44, 6074–6081.
- Grayver, A.V., Olsen, N., 2019. The magnetic signatures of the m_2 , n_2 , and o_1 oceanic tides observed in swarm and champ satellite magnetic data. *Geophysical Research Letters* 46, 4230–4238.
- Kowalik, Z., Proshutinsky, T., Proshutinsky, A., 2006. Tide-tsunami interactions. *Science of Tsunami Hazards* 24, 242–256.
- Maksimov, M., Velímský, J., 2017. Fast calculations of the gradient and the hessian in the time-domain global electromagnetic induction inverse problem. *Geophysical Journal International* 210, 270–283.
- Schmitz Jr, W.J., 1996. *On the World Ocean Circulation: Volume 2 The Pacific and Indian Oceans/A Global Update*. Technical Report. WOODS HOLE OCEANOGRAPHIC INSTITUTION MA.

- Thébault, E., Finlay, C.C., Beggan, C.D., Alken, P., Aubert, J., Barrois, O., Bertrand, F., Bondar, T., Boness, A., Brocco, L., et al., 2015. International geomagnetic reference field: the 12th generation. *Earth Planets Space* 67, 1–19.
- Tyler, R.H., Boyer, T.P., Minami, T., Zweng, M.M., Reagan, J.R., 2017. Electrical conductivity of the global ocean. *Earth Planets Space* 69, 156.
- Velímský, J., 2013. Determination of three-dimensional distribution of electrical conductivity in the Earth's mantle from Swarm satellite data: Time-domain approach. *Earth Planets Space* 65, 1239–1246.